

Study of a background reconstruction method for the measurement of $D - \overline{D}$ azimuthal angular correlations

Long Ma^{a,c,*}, Xin Dong^b, Yu-Gang Ma^{a,d}

^a*Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China*

^b*Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA*

^c*University of Chinese Academy of Sciences, Beijing 100049, China*

^d*Shanghai Tech University, Shanghai 200031, China*

Abstract

We study experimental background reconstruction methods for the measurement of $D - \overline{D}$ correlation using a PYTHIA simulation. Like-sign and side-band background methods that are widely used in the experimental measurements of single D -meson production yields were deployed for the correlation background study. It is found that the like-sign method which well describes the combinatorial background of single D^0 meson yields fails to reproduce the correlated background in the $D^0 - \overline{D}^0$ correlation measurement, while the side-band background method yields a good description of the background for both the single D^0 yields and of the correlated background of the $D^0 - \overline{D}^0$ correlation measurement.

Keywords: heavy flavor, correlation, PYTHIA

PACS:

1. Introduction

Quantum Chromodynamics (QCD) is the theory describing quarks, gluons and the strong interaction between them. In QCD, heavy flavor quarks (c , b) are mostly produced through initial hard scatterings in high energy collisions

*Corresponding author, email address: malong@sinap.ac.cn

of nucleons or nuclei. Because of their large masses, heavy quarks may offer a unique sensitivity to study the cold and hot QCD media created in these collisions [1–3]. In proton + proton ($p+p$) collisions, perturbative QCD (pQCD) calculations reproduce the inclusive heavy flavor hadron production cross section data over a broad range of collision energies and rapidities [4–8]. The nuclear modification factor for single charmed hadrons in heavy ion collisions shows a significant modification compared to the $p+p$ reference. Several models with different energy loss mechanisms can all describe the experimental data [9–13].

Recent research suggests that azimuthal correlations $\Delta\phi$ between heavy quark pairs offer new insight into charm-medium interaction dynamics, and therefore can help distinguish different energy loss mechanisms in the hot QCD medium [14, 15]. The theoretical prediction indicates that pure radiative energy loss does not change the initial angular correlation function significantly, while pure collisional energy loss is more efficient at diluting the initial back-to-back charm pair correlation. Furthermore, the momentum broadening in the direction perpendicular to the initial quark momentum, which cannot be probed directly with traditional single particle measurements (e.g. nuclear modification factor (R_{AA}) and flow parameter v_2), could be reflected in the azimuthal angle correlations [15, 16].

In $p + p$ collisions, charm quark pairs are produced through initial back-to-back hard scatterings in leading order. In next-to-leading order, the angular correlation between charm quark pairs is widening. In particular, it will show a near-side peak at $\Delta\phi \sim 0$ if the charm pairs are produced through gluon splitting. The measurement of $D - \bar{D}$ correlations in $p + p$ collisions not only provides a baseline for heavy ion measurements, but also offers a good constraint for pQCD calculations and helps us understand non-perturbative fragmentation and the underlying event activity distributions. D -mesons inherit most of the initial charm pair correlations, but weak decays smear the correlation significantly. Therefore measurement of $D - \bar{D}$ correlations should be the most sensitive probe to address charm quark pair correlations [17, 18].

The experimental reconstruction of $D - \bar{D}$ azimuthal angular correlation is

challenging. It requires reconstruction of two charmed hadrons in a single event. Charmed hadrons need to be reconstructed through their hadronic decay channels which typically have small branching ratios. Furthermore, there are often sizable uncorrelated and correlated background in each reconstructed charmed hadron. In single charmed hadron yield measurements, for instance D^0 mesons through the $K^-\pi^+$ decay channel, several background methods, e.g. like-sign (LS), side-band (SB) and mixed-event (ME) etc. were deployed by experimentalists. In the ME technique, background pairs are reconstructed using two daughter tracks from different events. Therefore this method has the advantage to reproduce the uncorrelated combinatorial background with good statistics, but it doesn't contain the correlated background part. In the LS technique, background is generated by pairing daughter tracks with the same charge sign. It contains also the correlated background assuming daughters are produced in pairs with opposite charge signs. In the SB technique, opposite sign pairs with invariant masses away from the D^0 peak are used, and usually two symmetric mass regions on both sides of the D^0 peak are selected and an average of these two is chosen to represent the background underneath the D^0 peak. To a reasonable precision, both LS and SB techniques can successfully reproduce the background in single D^0 yield measurements [19, 20].

In this paper, we investigate these background reconstruction methods for the measurement of $D - \bar{D}$ correlations. The ME technique misses the correlated background and it typically needs to be normalized to either LS or SB distributions. In the following study we focus on the comparison of the LS and SB background techniques.

2. PYTHIA study for $D - \bar{D}$ correlations

The Monte Carlo event generator PYTHIA (version 8.168) is used for this study [21]. We focus on $p + p$ collisions at $\sqrt{s_{\text{NN}}} = 500$ GeV for illustration. The parameters were tuned so that PYTHIA can reproduce the experimental data

on inclusive $c\bar{c}$ production cross-section in $p + p$ collisions at 500 GeV measured by the STAR experiment at RHIC [22].

Figure 1 shows the $c\bar{c}$ production cross section as a function of transverse momentum (p_T) in PYTHIA in comparison with the STAR measurements. The modified PYTHIA parameters in this tune are: strong interaction coupling constant (α_s) value of final parton shower (TimeShower:alphaSvalue) to be 0.15 and minimum invariant transverse momentum ($p_\perp$) threshold for hard QCD process (PhaseSpace:pTHatMin) to be 1.5 GeV/c. PYTHIA with this tune describes both the magnitude and the p_T spectrum well. It was also found that changing these two parameters has a negligible effect on charm correlations.

A sample of six billion PYTHIA minimum bias events with this tune were generated for the $D^0 - \bar{D}^0$ correlation study. D^0 mesons can be directly accessed in the PYTHIA simulation based on their particle identification number. To emulate the experimental measurement, D^0 s were reconstructed by pairing kaon and pion candidate pairs via the typical hadronic decay channel $D^0 \rightarrow K^- + \pi^+$ and its charge conjugate channel for $\bar{D}^0$. In a real experiment with a silicon vertex detector, many background daughter tracks from the primary collisions can be eliminated, but there remains considerable background, particularly in the low p_T region.

In this study, we don't distinguish secondary decay vertices in the D^0 reconstruction. Instead, we combine all kaons and pions at midrapidity (pseudorapidity $|\eta| \leq 1$) in the final stage in the PYTHIA output, which allows us to study the validity of the background reconstruction methods with different signal-to-background ratios in the reconstructed D^0 candidates. The invariant mass of unlike-sign and like-sign kaon and pion pairs in the same event are calculated. A finite momentum resolution effect is included so that the reconstructed D^0 signal peak has the width observed by the experiment.

Figure 2 demonstrates the $D^0(\bar{D}^0)$ signal and the combinatorial background from the like-sign method and the side-band method. The like-sign background and side-band background regions are denoted by the blue and red hatched areas, respectively. Invariant mass distribution of $\bar{D}^0$ is almost identical to D^0

in both shape and magnitude. The background is found to be flat in PYTHIA within a relative wide invariant mass range. For simplicity, we denote $D^0(\overline{D}^0)$ candidates from $K^-\pi^+(K^+\pi^-)$ pairs with unlike signs as ‘US’ candidates, and those from $K^-\pi^-$ or $K^+\pi^+$ pairs with same charge sign as ‘LS’ background. The side-band background is denoted as ‘SB’. Figure 2 shows both LS and SB methods can reasonably reproduce the real background underneath the reconstructed D^0 signals. For the single particle yield measurement, the D^0 and $\overline{D}^0$ counts are calculated by Eq. 1 and Eq. 2 for LS and SB background methods, respectively.




Figure 1: (Color online) Charm pair cross section as a function of transverse momentum in p + p collisions at $\sqrt{s_{NN}} = 500$ GeV in PYTHIA (dashed line) compared with STAR measurements (solid circles).




Figure 2: (Color online) Invariant mass distribution of all final stage kaon and pion pairs with opposite signs in PYTHIA data at mid-rapidity (shown by solid red line, US). Like-sign method reproduces the combinatorial background shown by the blue solid line (LS). The blue shaded region shows the like-sign background within a $\pm 3\sigma$ window of the signal peak. The side-band background regions are shaded red (SB).

If the background methods work well for the $D^0\text{-}\overline{D}^0$ correlation measurement, the correlation signal between D^0 and $\overline{D}^0$ can be derived using Eq. 3 and Eq. 4. The asterisks (*) indicate the correlation functions between the pairs.

We can also derive the $D^0\text{-}\overline{D}^0$ correlation signal from the PYTHIA simulation directly and compare to the reconstructed signals with these two background methods.

$$N_{LS}^{D^+\overline{D}^0} = US(K^-\pi^+) + US(K^+\pi^-) - LS(K^-\pi^-) - LS(K^+\pi^+) \quad (1)$$

$$N_{SB}^{D^+\overline{D}^0} = US(K^-\pi^+) + US(K^+\pi^-) - SB(K^-\pi^+) - SB(K^+\pi^-) \quad (2)$$

$$C_{LS}^{D\overline{D}} = US(K^-\pi^+) * US(K^+\pi^-) - LS(K^-\pi^-) * US(K^-\pi^+) \\ - LS(K^+\pi^+) * US(K^+\pi^-) + LS(K^-\pi^-) * LS(K^+\pi^+), \quad (3)$$

$$C_{SB}^{D\overline{D}} = US(K^-\pi^+) * US(K^+\pi^-) - SB(K^+\pi^-) * US(K^-\pi^+) \\ - SB(K^-\pi^+) * US(K^+\pi^-) + SB(K^-\pi^+) * SB(K^+\pi^-), \quad (4)$$




Figure 3: (Color online) Left figure: cross correlations of $D^0 - \overline{D}^0$ from unlike-sign candidates(US) and like-sign backgrounds(LS). The trigger and associate p_T cuts are both set to 1 GeV/c with a signal-to-background ratio of around 0.3. The right figure shows similar results from the side-band method.

The di-hadron correlation measurements are usually plotted as a function of the azimuthal angle difference, i.e. $\Delta\phi = \phi_{D^0} - \phi_{\overline{D}^0}$. In Figure 3, the left plot shows the correlations between unlike-sign candidates and the like-sign

backgrounds as a function of $\Delta\phi$. The $p_T > 1.0 \text{ GeV}/c$ cut is set for both D^0 and $\overline{D^0}$ mesons, and the mass window cuts for US, LS and SB pairs used are shown as the colored bands in Fig. 2. The plot shows that the correlation between like-sign and like-sign background pairs (LS*LS) tends to peak at $\Delta\phi$ around 0 and that its magnitude is considerably larger than that between like-sign background and unlike-sign candidates (LS*US). The right panel in Figure 3 shows the results with the same trigger p_T and S/B ratio from the side-band method. The correlation between the side-band background and unlike-sign candidates lies between the other two correlation terms. The correlation between two side-band background pairs shows a similar trend as that between side-band background and unlike-sign candidates.




Figure 4: (color online). The $D^0 - \overline{D^0}$ correlation as a function of relative azimuthal angle $\Delta\phi$ in p+p collisions at $\sqrt{s_{NN}} = 500 \text{ GeV}$ calculated with like-sign method (left) and side-band method(right) based on Equation. 3 in PYTHIA simulation. The transverse momentum dependence is shown with p_T cuts applied to triggered and associated D mesons. Panels (a)-(d) show correlations of reconstructed D^0 mesons under different S/B ratios in comparison with correlations of produced $D^0 - \overline{D^0}$ pairs in PYTHIA.

In Fig. 4, the reconstructed $D^0 - \overline{D^0}$ correlation signals with LS and SB background methods are compared to the real correlation signals from PYTHIA directly. Two sets of p_T cuts were imposed for triggered and associated D -mesons, shown in the upper and lower panels respectively. Panels in two different

columns show the comparisons with two different mass window cuts which result in different signal-to-background ratios in the reconstructed D^0 candidates. The red data points represent the correlation signals from reconstructed D^0 s, and the blue data points are the real D^0 correlations directly obtained from PYTHIA with the same kinematic cuts applied. Similar results from the SB method are also shown in Fig. 4.

As we can see, reconstructed correlations using the LS method are different from the real $D^0\text{-}\overline{D}^0$ signal from PYTHIA. Particularly, the reconstruction correlations start to show an enhanced structure in the near side region when the signal-to-background ratio is getting smaller. While reconstructed correlations using the SB method can reproduce the real $D^0\text{-}\overline{D}^0$ signal reasonably well in these kinematic and signal-to-background ratio regions. It is also found that the quality of the agreement does not depend on the transverse momentum cut. It depends on the signal-to-background ratios of the D^0 candidates.

$$u_i = \left(\frac{1}{N_{trig}} \frac{dN}{d\Delta\phi_i} \right)_{reco}, \quad v_i = \left(\frac{1}{N_{trig}} \frac{dN}{d\Delta\phi_i} \right)_{real} \quad (5)$$

$$\Delta P = \frac{1}{N} \sum_{i=1}^k \left| \frac{u_i - v_i}{v_i} \right|, \quad \Delta E = \frac{1}{N} \sum_{i=1}^k \left| \frac{\sqrt{\sigma_{u_i}^2 + \sigma_{v_i}^2}}{v_i} \right| \quad (6)$$

To better illustrate the performance of these two background methods in measuring the angular correlations of the $D - \overline{D}$ pairs, we define two variables to quantify the goodness of fit for the reconstructed correlation signals with respect to the real $D^0 - \overline{D}^0$ correlations from PYTHIA. ΔP and ΔE are defined in Equation. 6 to describe the relative difference of the data points and the statistical errors from this sample. u_i and v_i are values of the number i data points of reconstructed and real correlation signals in each $\Delta\phi$ bin. N is the total number of data points in each correlation signal assuming the same binning for the histograms. Fig. 5 shows the corresponding results from like-sign method and side-band methods respectively. As we can see, ΔP in like-sign results shows a large deviation from ΔE when the S/B ratio goes down indicating that like-sign method fails to reproduced real correlation at relatively low S/B ratios.




Figure 5: (color online). The summary plots of the goodness of fit calculated with like-sign method (left) and side-band method (right) in PYTHIA simulation. The estimator is shown as a function of the signal over background ratio (S/B). Solid line and dashed line show ΔP and ΔE respectively.

While the side-band method shows a good performance throughout the whole S/B ratio region investigated. The increase of both ΔP and ΔE at low S/B region in the side-band method is due to the reduced statistics.

When a $K^+\pi^+$ pair is selected, there is a higher probability for finding a $K^-\pi^-$ pair than another $K^+\pi^+$ pair due to local and global charge conservation. The reconstructed correlation signal after like-sign background subtraction from Eq. 3 should contain all correlations between $K^+\pi^-$ and $K^-\pi^-$ pairs including the $D^0 - \overline{D^0}$ correlation we are interested in as well as the correlation due to charge conservation. To further demonstrate that the additional correlation observed in the like-sign method is related to the underlying event instead of the $D^0 - \overline{D^0}$ signal, we turn off D^0 hadronic decay process in the PYTHIA simulation and run the same analysis.

Fig. 6 shows the invariant mass distribution of pure $K\pi$ pairs without D^0 decay contribution. Cross correlations between unlike-sign/like-sign, like-sign/like-sign pairs are plotted in comparison with the unlike-sign/unlike-sign pair correlations in Fig. 7 with different cuts applied to the invariant mass region. Similarly, results from the side-band method are shown in the right panel




Figure 6: (Color online) Invariant mass distribution of pure $K\pi$ pairs in PYTHIA. $D^0 \rightarrow K\pi$ hadronic decay process is turned off. Black line: Unlike-sign $K\pi$ pairs. Black hatched area: Like-sign $K\pi$ pairs within cut window. Blue shaded area: Side-band $K\pi$ pairs within cut window.

of Fig. 7. There is a large difference between the LS*US pair correlation and the US*US pair correlation while there is very small difference between LS*LS and US*US. This is consistent with our understanding that there is an additional correlation not originating from the $D^0 - \overline{D}^0$ pairs.

The side-band method is not affected by charge conservation. It is shown that all cross correlations fall in the same trend, and there is no remaining $K^+\pi^- - K^-\pi^+$ correlation when the $D^0 \rightarrow K^+\pi^-$ decay is turned off.

3. Conclusion

In summary, we study the background reconstruction methods for azimuthal correlations between D^0 and $\overline{D}^0$ pairs by a PYTHIA simulation.

Both like-sign and side-band methods show a good description of the background when reconstructing single D^0 yields. However, when reconstructing the correlation signal, the like-sign method fails to reproduce the $D^0 - \overline{D}^0$ correlation. The residual correlation after like-sign background subtraction mainly




Figure 7: (Color online) Left Panel: Cross correlations of the pure like-sign(LS) and unlike-sign(US) $K\pi$ pairs in like-sign method. Right Panel: Cross correlations of side-band(SB) and unlike-sign(US) $K\pi$ pairs in side-band method.

comes from the underlying event activity likely due to local and/or global charge conservation. We demonstrate that the side-band method shows a good performance in describing the correlation background and therefore reproduce the original $D^0 - \overline{D}^0$ correlation in the signal-to-background ration regions investigated.

4. Acknowledgement

We thank Dr. Hans Georg Ritter for the careful English language correction. This work was supported in part by the National Natural Science Foundation of China under contract Nos. 11421505 and the Major State Basic Research Development Program in China under Grant No. 2014CB845400.

References

References

- [1] Rudolph C. Hwa, Phys. Rev. Lett. 77, 2 (1996)
- [2] B. Svetitsky, A. Uziel, Phys. Rev. D 55, 2616 (1997)
- [3] Z. Lin and M. Gyulassy, Phys. Rev. C 51, 2177 (1995)
- [4] C. Albajar et al. (UA1 Collaboration), Phys. Lett. B 244, 566 (1990)
- [5] F. Abe et al. (CDF Collaboration), Phys. Rev. Lett. 68, 3403 (1992)
- [6] D. Acosta et al. (CDF Collaboration), Phys. Rev. Lett. 91, 241804 (2003)
- [7] B. Abelev et al. (ALICE Collaboration), JHEP 01, 128 (2012)
- [8] B. Abelev et al. (ALICE Collaboration), JHEP 1207, 191 (2012)
- [9] M. Cacciari, M. Greco, and P. Nason, JHEP 9805, 007 (1998)
- [10] M. Cacciari, S. Frixione, and P. Nason, JHEP 0103, 006 (2001)
- [11] T. Sjostrand, S. Mrenna, and P. Skands, JHEP 0605, 026 (2006)
- [12] Yi-Fei Xu, et al., Nuclear Science and Techniques, 27, 87 (2016)
- [13] Jie Chen, et al., Nuclear Science and Techniques, 27, 73 (2016)
- [14] S. Cao, G.Y. Qin and S. A. Bass, arXiv:1404.1081.
- [15] M. Nahrgang, et al., Phys. Rev. C 90, 024907 (2014)
- [16] S. Cao, G.Y. Qin and S. A. Bass, Phys. Rev. C 92, 054909 (2015)
- [17] X. Zhu, et al., Phys. Lett. B 647, 152301 (2007)
- [18] X. Zhu, N. Xu and P. Zhuang, Phys. Rev. Lett. 100, 152301 (2008)
- [19] L. Adamczyk, et al., (STAR Collaboration), Phys. Rev. D 86, 072013 (2012).

- [20] L. Adamczyk, et al., (STAR Collaboration), Phys. Rev. Lett. 113, 142301 (2014).
- [21] Torbjörn Sjöstrand, Stephen Mrenna, Peter Skands, Comput.Phys.Commun.178:852-867 (2008)
- [22] David Tlusty (for the STAR collaboration) Journal of Physics: Conference Series 509, 012078 (2014)